

PHYTOCHEMICALS AND BIOACTIVITIES OF *ARTOCARPUS*  
*LANCEIFOLIUS* ROXB AND *ARTOCARPUS* MAINGAYI KING

NOOR SAFINA SULAIMAN

UNIVERSITI TEKNOLOGI MALAYSIA

**PHYTOCHEMICALS AND BIOACTIVITIES OF *ARTOCARPUS*  
*LANCEIFOLIUS* ROXB AND *ARTOCARPUS* MAINGAYI KING**

**NOOR SAFINA SULAIMAN**

**A thesis submitted in fulfillment of the  
requirements for the award of the degree of  
Master of Science (Chemistry)**

**Faculty of Science  
Universiti Teknologi Malaysia**

**JULY 2011**

*Dedicated to*

*My beloved parents*

*My sisters and brother*

*My friends.*

## ACKNOWLEDGEMENT

I thank to Allah S.W.T for His love and care which kept me going forward and for His grace that led me throughout the whole process of completing this research. I praise His faithfulness.

I would like to express my sincere gratitude, appreciation and thank to my supervisor, Dr. Shajarahtunnur Jamil, for all her guidance, continuous support and her patience throughout the completion of this research study.

Special thanks also goes to Assoc. Prof. Dr. Farediah Ahmad, Professor Dr. Hasnah Mohd Sirat, and Assoc. Prof. Dr. Zakaria Bahari who kept an eye on the progress of my work, for their advice and moral support.

I also would like to thank to E-science Fund for granting me scholarships. My sincere appreciation also extends to laboratory assistants who help me in using several instruments in the Department of Chemistry, Faculty of Science.

I would also like to thank all my colleagues and friends especially Ms Nadya, Ms. Della, Ms. Noraini, Ms. Athirah, Ms. Raihan, Mr. Nuzul, Mr. Tarmizi, Ms. Mariami Alisa, and Ms. Susilawati for their help, discussions and encouragement.

Finally I would express my deepest appreciations to my loving parents for all their love, sacrifice, understanding and support. Without them, I could never be succeeded.

## PREFACE

This thesis is the resulted of my own work carried out in the Department of Chemistry, Faculty of Science, Universiti Teknologi Malaysia between June 2008 and July 2010 under the supervision of Dr. Shajarahtunnur Jamil. Part of my work describe in this thesis has been reported in the following publications:

1. Noor Safina Sulaiman and Shajarahtunnur Jamil (2009). "Screening of *Artocarpus lanceifolius* for Antibacterial Activity". Second International Conference and Workshops & Regional Fundamental Science Seminar 2009, 2-4 June 2009, Johor Bahru, Malaysia, **1**, 242-244, ISBN : 978-983-9805-73-4
2. Noor Safina Sulaiman, Shajarahtunnur Jamil (2009). "Phytochemicals Studies and Antimicrobial Activities of *Artocarpus lanceifolius*". 2<sup>nd</sup> Junior Chemist Colloquium 2009, 1-2 July 2009, UNIMAS Sarawak, Malaysia.

## ABSTRACT

Phytochemical studies on *Artocarpus lanceifolius* Roxb. and *Artocarpus maingayi* King have successfully isolated seven flavonoids and two plant sterols. Five compounds were successfully isolated from the stem bark of *Artocarpus lanceifolius* which were identified as 9,19-cyclolanost-24-en-3-acetate, cycloartobiloxanthone, artonol B, 7,8-(2,2-dimethylchromeno)-10-(2-hydroxyl-1-methylethyl)-2',4',5',5-tetrahydroxyflavone, and artonin E. Two flavans which were tentatively elucidated as 7,8-(2,2-dimethylchromano)-4',4,5-trihydroxy-2'-methoxyflavan and 8-(11-methyl-11-butenyl)-3,4',5,7-tetrahydroxy-2'-methoxyflavan have been isolated from the leaves of *Artocarpus lanceifolius*. Another two compounds were isolated from the leaves of *Artocarpus maingayi* which were identified as carpachromene, and  $\beta$ -sitosterol. Structures of all compounds were elucidated spectroscopically by Nuclear Magnetic Resonance, Infrared, Ultraviolet Spectroscopies and Mass Spectrometry. Biological activities were carried out on the crude extracts and pure compounds. The antimicrobial testing on the crude extracts and pure compounds were carried out against the Gram-positive bacteria, *Bacillus subtilis* and *Staphylococcus aureus* and Gram-negative bacteria, *Escherichia coli* and *Pseudomonas aeruginosa*. Most of the crude extracts and pure compounds showed significant antimicrobial activity. A pure compound, 7,8-(2, 2-Dimethylchromeno)-10-(2-hydroxyl-1-methylethyl)-2',4',5',5-tetrahydroxyflavone showed most significant antimicrobial activity compared to other isolated compounds and crude extracts with the minimum bactericidal concentration and minimum inhibition concentration value between 31.25 – 62.5  $\mu\text{g/mL}$ . The antioxidant test on the crude extracts and pure compounds from both plant species showed potential free radical scavenger against 2,2-diphenyl-1-picrylhydrazyl. Artonin E exhibited the strongest antioxidant activity with 50% scavenging concentration at 46.9  $\mu\text{g/mL}$  while the ethyl acetate crude extract from the stem bark of *Artocarpus maingayi* showed 50% scavenging concentration at 60.1  $\mu\text{g/mL}$ .

## ABSTRAK

Kajian fitokimia ke atas *Artocarpus lanceifolius* Roxb. dan *Artocarpus maingayi* King berjaya mengasingkan tujuh sebatian flavonoid dan dua sebatian sterol tumbuhan. Lima sebatian telah berjaya diasingkan daripada kulit batang *Artocarpus lanceifolius* yang di kenalpasti sebagai 9,19-siklolanos-24-en-3-asetat, sikloartobiloxanton, artonol B, 7,8-(2,2-dimetilkromeno)-10-(2-hidroksil-1-metiletil)-2',4',5',5-tetrahidroksiflavan, dan artonin E. Dua flavan yang dinamakan sebagai 7,8-(2,2-dimetilkromano)-4',4,5-trihidroksi-2'-metoksiflavan dan 8-(11-metil-11-butenil)-3,4',5,7-tetrahidroksi-2'-metoksiflavan telah diasingkan daripada daun *Artocarpus lanceifolius*. Dua lagi sebatian kimia telah diasingkan daripada daun *Artocarpus maingayi* yang dikenalpasti sebagai karpakromen dan  $\beta$ -sitosterol. Struktur semua sebatian kimia ini dikenalpasti berdasarkan kepada kajian spektroskopi resonans magnet nukleus, infra merah, spektroskopi ultralembayung dan spektrometri jisim. Kajian aktiviti biologi telah dijalankan ke atas ekstrak mentah dan sebatian tulen. Ujian antibakteria ke atas ekstrak mentah dan sebatian tulen telah dijalankan dengan menggunakan Gram positif, *Bacillus subtilis* dan *Staphylococcus aureus* manakala Gram-negatif, *Escherichia coli* dan *Pseudomonas aeruginosa*. Kebanyakan ekstrak mentah dan sebatian tulen menunjukkan aktiviti antibakteria yang signifikan. Satu sebatian tulen, 7,8-(2,2-dimetilkromeno)-10-(2-hidroksil-1-metiletil)-2',4',5',5-tetrahidroksiflavan menunjukkan aktiviti antibakteria yang paling signifikan berbanding dengan sebatian tulen yang lain dengan nilai kepekatan minimum bakterisidal dan kepekatan minimum perencatan di antara 31.25 hingga 62.5  $\mu\text{g/mL}$ . Ujian antioksidan ke atas ekstrak mentah dan sebatian tulen dari kedua-dua spesies *Artocarpus* menunjukkan keupayaan untuk bertindak sebagai perencat radikal bebas terhadap 2,2-difenil-1-picrylhidrazil. Artonin E menunjukkan aktiviti antioksidan yang terkuat dengan nilai kepekatan untuk merencat 50% radikal pada 46.9  $\mu\text{g/mL}$ , manakala ekstrak mentah etil asetat daripada kulit batang *Artocarpus maingayi* menunjukkan nilai kepekatan untuk merencat 50% radikal pada 60.1  $\mu\text{g/mL}$ .